



LAWRENCE  
LIVERMORE  
NATIONAL  
LABORATORY

# Consequences of the Large-Scale Subsidence Rate on the Stably Stratified Atmospheric Boundary Layer Over the Arctic Ocean, as seen in Large-Eddy Simulations

J. D. Mirocha, B. Kosovic

January 19, 2006

American Meteorological Society's 17th Symposium on  
Boundary Layers and Turbulence  
San Diego, CA, United States  
May 22, 2006 through May 26, 2006

## **Disclaimer**

---

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

**Consequences of the Large-Scale Subsidence Rate on the Stably Stratified  
Atmospheric Boundary Layer Over the Arctic Ocean, as  
seen in Large-Eddy Simulations**

Abstract for the American Meteorological Society's  
**17th Symposium on Boundary Layers and Turbulence**  
Monday, 22 May 2006 - Friday, 26 May 2006  
San Diego, CA

Jeffrey D. Mirocha  
Branko Kosovic  
Lawrence Livermore National Laboratory  
Livermore, CA 94551  
(925) 422-4627, [mirocha2@llnl.gov](mailto:mirocha2@llnl.gov)

**Abstract**

The analysis of surface heat fluxes and sounding profiles from SHEBA indicated possible significant effects of subsidence on the structure of stably-stratified ABLs (Mirocha et al. 2005). In this study the influence of the large-scale subsidence rate on the stably stratified atmospheric boundary layer (ABL) over the Arctic Ocean during clear-sky, winter conditions is investigated using a large-eddy simulation model. Simulations are conducted while varying the subsidence rate between 0, 0.001 and 0.002  $\text{ms}^{-1}$ , and the resulting quasi-equilibrium ABL structure and evolution are examined. Simulations conducted without subsidence yield ABLs that are deeper, more strongly mixed, and cool much more rapidly than were observed. The addition of a small subsidence rate significantly improves agreement between the simulations and observations regarding the ABL height, potential temperature profiles and bulk heating rates. Subsidence likewise alters the shapes of the surface-layer flux, stress and shear profiles, resulting in increased vertical transport of heat while decreasing vertical momentum transport. A brief discussion of the relevance of these results to parameterization of the stable ABL under subsiding conditions in large-scale numerical weather and climate prediction models is presented.

This work was performed under the auspices of the U.S. Department of Energy by University of California, Lawrence Livermore National Laboratory under Contract W-7405-Eng-48.